

Radiation Fluid Stars in the Non-minimally Coupled $Y(R)F^2$ Gravity

Özcan SERT*

*Department of Mathematics, Faculty of Arts and Sciences,
Pamukkale University, 20070 Denizli, Türkiye*

(Dated: January 31, 2017)

Abstract

We propose a non-minimally coupled gravity model in $Y(R)F^2$ form to describe the radiation fluid stars which have the radiative equation of state between the energy density ρ and the pressure p as $\rho = 3p$. Here F^2 is the Maxwell invariant and $Y(R)$ is a function of the Ricci scalar R . We give the gravitational and electromagnetic field equations in differential form notation taking the infinitesimal variations of the model. We look for electrically charged star solutions to the field equations under a constraint which eliminating complexity of the higher order terms in the field equations. We determine the non-minimally coupled function $Y(R)$ and the corresponding model which admits new exact solutions in the interior of star and Reissner-Nordstrom solution at the exterior region. Using vanishing pressure condition at the boundary together with the continuity conditions of the metric functions and the electric charge, we find the mass-radius ratio, charge-radius ratio and gravitational surface redshift depending on the parameter of the model for the radiation fluid star. We derive general restrictions for the ratios and redshift of the charged compact stars. We obtain a slightly smaller upper mass-radius ratio limit than the Buchdahl bound $4/9$ and a smaller upper redshift limit than the bound of the standard General Relativistic stars.

* osert@pau.edu.tr

I. INTRODUCTION

The radiation fluid stars have crucial importance in astrophysics. They can describe the core of neutron stars which is a collection of cold degenerate (non-interacting) fermions [1–4] and self-gravitating photon stars [4–8]. Such radiative stars which called Radiation Pressure Supported Stars (RPSS) can be possible even in Newtonian Gravity[9] and their relativistic extension which called Relativistic Radiation Pressure Supported Stars (RRPSS) [10] can describe the gravitational collapse of massive matter clouds to very high density fluid. There are also some investigations related with gedanken experiments such as black hole formation and evaporation with self-gravitating gas confined by a spherical symmetric box. These investigations [11, 12] can lead to new insights to the nature of Quantum Gravity.

The radiation fluid stars have the radiative equation of state with $\rho = 3p$ which is the high density limit of the general isothermal spheres satisfying the linear barotropic equation of state $\rho = kp$ with constant k . The entropy and thermodynamic stability of self-gravitating charge-less radiation fluid stars were firstly calculated in [5] using Einstein equations. This work was extended to the investigation of structure, stability and thermodynamic parameters of the isothermal spheres involving photon stars and the core of neutron stars [7, 8]. Also, the numerical study of such a charge-less radiative star which consists of a photon gas conglomerations can be found in [6]. Some interesting interior solutions of the General Relativistic field equations in isotropic coordinates with the linear barotropic equation of state were presented by Mak and Harko [13–15] for dense astrophysical objects without charge.

Furthermore, a spherically symmetric fluid sphere which contains a constant surface charge can be more stable than the charge-less case [16]. The gravitational collapse of a spherically symmetric star may be prevented by charge [17], since the repulsive electric force contributes to counterpoise the gravitational attraction [18]. It is interesting to note that interior of a strange quark star can be described by a charged solution admitting a one-parameter group of conformal motions [19] for the equation of state $\rho = 3p + 4B$ which is known as the MIT bag model. The physical properties and structure of the radiation fluid stars in the model with hybrid metric-Palatini gravity [20] and Eddington-inspired Born-Infeld (EIBI) gravity [21] were obtained numerically. It is a challenging problem to find exact interior solutions

of the the charged radiation fluid stars, since the trace of the gravitational field equations gives zero Ricci scalar for the radiative equation of state $\rho = 3p$. Therefore it is important to find a modified gravity model which can describe the radiation fluid stars analytically.

In this study we propose a non-minimally coupled modified gravity model in $Y(R)F^2$ -form in order to find exact solutions to the radiation fluid stars. Here F^2 is the Maxwell invariant and $Y(R)$ is a function of the Ricci scalar R . We will determine the non-minimal function from physically applicable solutions of the field equations and boundary conditions. Such a coupling in RF^2 form was first introduced by Prasanna [22] to understand the intricate nature between all energy forms, electromagnetic fields and curvature. Later, a class of such couplings was investigated to gain more insight on charge conservation and curvature [23]. These non-minimal terms can be obtained from dimensional reduction of a five dimensional Gauss-Bonnet gravity action [24] and R^2 -type action [25, 26]. The calculation of QED photon propagation in a curved background metric [27] leads to these terms. A generalization of the non-minimal model to $R^n F^2$ -type couplings [28–33] may explain the generation of seed magnetic fields during the inflation and the origin of large-scale magnetic fields in the universe [28–30]. Another generalization of the non-minimal RF^2 model to non-Riemannian space-times [34] can give more insights to the torsion and electromagnetic fields. Then it is possible to consider the more general couplings with any function of the Ricci scalar and the electromagnetic fields such as $Y(R)F^2$ -form. These non-minimal models in $Y(R)F^2$ -form have very interesting solutions such as regular black hole solutions to avoid singularity [35], spherically symmetric static solutions to explain the rotation curves of galaxies [33, 36–38], cosmological solutions to explain cosmic acceleration of the universe [32, 39–41] and pp-wave solutions [42].

In order to investigate a charged astrophysical phenomena one can consider Einstein-Maxwell theory which is a minimal coupling between gravitational and electromagnetic fields. But when the astrophysical phenomena has high density, pressure and charge such as neutron stars and quark stars, new interaction types between gravitational and electromagnetic fields may be appeared. Then the non-minimally coupled $Y(R)F^2$ Gravity can be ascribed to such new interactions and we can apply the theory to the charged compact stellar system. In this study we focus on exact solutions of the radiation fluid stars for the non-minimally coupled model, inspired by the solution in [19]. We construct the non-

minimal coupling function $Y(R)$ with the parameter α and the corresponding model. We give interior and exterior solutions of the model. Similarly to [19], our interior solutions turn out to the solution given by Misner and Zapsky [2] with $b = 0$ and $Q = 0$, describing ultra high density neutron star or the relativistic Fermi gas. We determine the total mass, charge and surface gravitational redshift of the stars depending on the boundary radius r_b and the parameter α using the matching conditions. We give the general restrictions for the ratios and redshift of the charged compact stars and compare them with the bound given in [43] and the Buchdahl bound [44].

The organization of the present work is as follows: The general action in $Y(R)F^2$ form and the corresponding field equations are given in Section II to describe a charged compact star. The spherically symmetric, static exact solutions under conformal symmetry and structure of the non-minimal function $Y(R)$ are obtained in Section III. Using the continuity and boundary conditions, the gravitational mass, total charge and redshift of the star are derived in Section IV. The conclusions are given in the last Section.

II. THE MODEL WITH $Y(R)F^2$ -TYPE COUPLING FOR A COMPACT STAR

The recent astronomical observations about the problems such as dark matter [45, 46] and dark energy [47–52] strongly support that Einstein’s theory of gravity needs to modification at large scales. Therefore Einstein-Maxwell theory also may be modified [28–30, 32, 33, 36–41] to explain these observations. Furthermore, since such astrophysical phenomenons as neutron stars or quark stars have high energy density, pressure and electromagnetic fields, new interaction types between gravitational and electromagnetic fields in $Y(R)F^2$ form can be appeared. When the extreme conditions are removed, this model turns out to the minimal Einstein-Maxwell theory. We write the following action to describe the interior of a charged compact star by adding the matter part L_{mat} and the source term $A \wedge J$ to the $Y(R)F^2$ -type non-minimally coupled model in [33, 35–41],

$$I = \int_M \left\{ \frac{1}{2\kappa^2} R * 1 - Y(R) F \wedge * F + 2A \wedge J + L_{mat} + \lambda_a \wedge T^a \right\} \quad (1)$$

depending on the fundamental variables such that the co-frame 1-form $\{e^a\}$, the connection 1-form $\{\omega^a_b\}$ and the homogeneous electromagnetic field 2-form F . We derive F from the

electromagnetic potential 1-form A such as $F = dA$. We constrain the model to the case with zero torsion connection by λ_a Lagrange multiplier 2-form. Then the variation of λ_a leads to the Levi-Civita connection which can be found from $T^a = de^a + \omega^a_b \wedge e^b = 0$. In action (1), J is the electric current density 3-form for the source fluid inside the star, $Y(R)$ is any function of the curvature scalar R . The scalar can be derived from the curvature tensor 2-forms $R^a_b = d\omega^a_b + \omega^a_c \wedge \omega^c_b$ via the interior product ι_a such as $\iota_b \iota_a R^{ab} = R$. We denote the space-time metric by $g = \eta_{ab}e^a \otimes e^b$ which has the signature $(-+++)$. Then we set the volume element with $*1 = e^0 \wedge e^1 \wedge e^2 \wedge e^3$ on the four dimensional manifold.

For a charged isotropic perfect fluid, Electromagnetic and Gravitational field equations of the non-minimal model are found from infinitesimal variations of the action (1)

$$d(*YF) = J, \quad (2)$$

$$dF = 0, \quad (3)$$

$$\begin{aligned} -\frac{1}{2\kappa^2}R_{bc} \wedge *e^{abc} &= Y(\iota^a F \wedge *F - F \wedge \iota^a *F) + Y_R F_{mn} F^{mn} *R^a \\ &+ D[\iota_b d(Y_R F_{mn} F^{mn})] \wedge *e^{ab} + (\rho + p)u^a *u + p *e^a, \end{aligned} \quad (4)$$

where $Y_R = \frac{dY}{dR}$ and $u = u_a e^a$ is the velocity 1-form associated with an inertial time-like observer, $u_a u^a = -1$. The modified Maxwell equation (2) can be written as

$$d * \mathcal{G} = J \quad (5)$$

where $\mathcal{G} = YF$ is the excitation 2-form in the interior medium of the star. More detailed analysis on this subject can be found in [42, 53, 54]. Following Ref. [40] we write the gravitational field equations (4) as follows:

$$G^a = \kappa^2 \tau_N^a + \kappa^2 \tau_{mat}^a, \quad (6)$$

where G^a is Einstein tensor $G^a = -\frac{1}{2}R_{bc} \wedge *e^{abc}$, τ_N^a and τ_{mat}^a are two separate effective energy momentum tensors, namely, the energy momentum tensor of the non-minimally coupled term introduced in [35, 40] and energy momentum tensor of matter, respectively

$$\tau_N^a = Y(\iota^a F \wedge *F - F \wedge \iota^a *F) + Y_R F_{mn} F^{mn} *R^a$$

$$+D[\iota_b d(Y_R F_{mn} F^{mn})] \wedge *e^{ab} , \quad (7)$$

$$\tau_{mat}^a = \frac{\delta L_{mat}}{\delta e_a} = (\rho + p)u^a * u + p * e^a . \quad (8)$$

We take the exterior covariant derivative of the modified gravitational equation (6) in order to show that the conservation of the total energy-momentum tensor $\tau^a = \tau_N^a + \tau_{mat}^a$

$$DG^a = D\tau^a . \quad (9)$$

The left hand side of equation (9) is identically zero $DG^a = 0$. The right hand side of equation (9) is calculated term by term as follows:

$$D[Y(\iota^a F \wedge *F - F \wedge \iota^a *F)] = 2J \wedge F^a - \frac{1}{2}F_{mn}F^{mn}dY \wedge *e^a \quad (10)$$

$$D(Y_R F_{mn} F^{mn} * R^a) = d(Y_R F_{mn} F^{mn}) \wedge *R^a + \frac{1}{2}F_{mn}F^{mn}Y_R dR \wedge *e^a \quad (11)$$

$$D[D[\iota_b d(Y_R F_{mn} F^{mn})] \wedge *e^{ab}] = D^2[\iota_b d(Y_R F_{mn} F^{mn})] \wedge *e^{ab} \quad (12)$$

$$= R_b^c \wedge \iota_c d(Y_R F_{mn} F^{mn}) * e^{ab} \quad (13)$$

$$= -d(Y_R F_{mn} F^{mn}) \wedge *R^a . \quad (14)$$

If we substitute all the expressions in (9) we find

$$0 = D\tau^a = 2J \wedge F^a + D\tau_{mat}^a \quad (15)$$

which leads to

$$D\tau_{mat}^a = -2J \wedge F^a \quad (16)$$

which is similar to the minimally coupled Einstein-Maxwell theory, but where $J = d(*YF)$ from (2). Then in this case without source $J = 0$, the conservation of energy-momentum tensor becomes $D\tau^a = 0 = D\tau_{mat}^a$.

The isotropic matter has the following energy density and pressure $\rho = \tau_{mat}^{0,0}$, $p = \tau_{mat}^{1,1} = \tau_{mat}^{2,2} = \tau_{mat}^{3,3}$ as the diagonal components of the matter energy momentum tensor τ_{mat}^a in the interior of the star. In order to get over higher order derivatives and the complexity of the

last term in (7) we take the following constraint

$$Y_R F_{mn} F^{mn} = \frac{K}{\kappa^2} \quad (17)$$

where K is a non-zero constant. If one take K is zero, then the non-minimal function Y becomes a constant and this is not different from the well known minimally coupled Einstein-Maxwell theory. The constraint (17) has the following futures: First of all, this constraint (17) is not an independent equation from the field equations, since the exterior covariant derivative of the gravitational field equations under the condition gives the constraint again in addition to the conservation equation. Secondly, the field equations (2-4) under the condition (17) with $K = -1$, can be interpreted [40] as the field equations of the trace-free Einstein gravity [55, 56] or unimodular gravity [57, 58] which coupled to electromagnetic energy-momentum tensor with the non-minimal function $Y(R)$, which are viable for astrophysical and cosmological applications. Thirdly, the constraint allows us to find the other physically interesting solutions of the non-minimal model [32, 33, 35, 36, 38, 40, 41]. Fourthly, when we take the trace of the gravitational field equation (4) as done in Ref. [40], we obtain

$$\frac{K+1}{\kappa^2} R * 1 = (\rho - 3p) * 1. \quad (18)$$

We can consider two cases satisfying (18) for the non-minimal $Y(R)F^2$ coupled model:

1. $K = -1$ which leads to the equation of state $\rho = 3p$ for the radiation fluid stars.
2. $K \neq -1$ with the equation $R = \frac{\kappa^2(\rho-3p)}{K+1}$.

Then we set $K = -1$ in (17) and (18), since we concentrate on the radiation fluid star for the non-minimal model. Therefore we see that the trace of the gravitational field equations does not give a new independent equation as another feature of the condition (17) with $K = -1$. One may see Ref. [40] for a detailed discussion on the physical properties and features of τ_N^a for the case $K = -1$. We leave the second case with $K \neq -1$, $\rho \neq 3p$ for next studies. We also note that the non-minimally coupled $Y(R)F^2$ model does not give any new solution for the MIT bag model $\rho - 3p = 4B$ with $B \neq 0$, since the curvature scalar R becomes a constant in (18) therewith $Y(R)$ must be constant. Thus this case is not a new model but the minimal Einstein-Maxwell case.

III. STATIC, SPHERICALLY SYMMETRIC CHARGED SOLUTIONS

We seek solutions to the model with $Y(R)F^2$ -type coupling describing a radiative compact star for the following most general (1+3)-dimensional spherically symmetric, static metric

$$ds^2 = -f^2(r)dt^2 + g^2(r)dr^2 + r^2 d\theta^2 + r^2 \sin(\theta)^2 d\phi^2 \quad (19)$$

and the following electromagnetic tensor 2-form with the electric field component $E(r)$

$$F = E(r)e^1 \wedge e^0. \quad (20)$$

We take the electric current density as a source of the field which has only the electric charge density component $\rho_e(r)$

$$J = \rho_e(r)e^1 \wedge e^2 \wedge e^3 = \rho_e g r^2 \sin \theta dr \wedge d\theta \wedge d\phi. \quad (21)$$

Using Stokes theorem, integral form of the Maxwell equation (2) can be written as

$$\int_V d * YF = \int_{\partial V} * YF = \int_V J = 4\pi q(r) \quad (22)$$

over the sphere which has the volume V and the boundary ∂V . When we take the integral, we find the charge inside the volume with the radius r

$$YEr^2 = q(r) = \int_0^r \rho_e(x)g(x)x^2 dx. \quad (23)$$

In (23), the second equality says that the electric charge can also be obtained from the charge density $\rho_e(r)$ of the star. Then the Gravitational field equations (4) lead to the following differential equations for the metric (19) and electromagnetic field (20) of the radiation fluid star $\rho = 3p$

$$\frac{1}{\kappa^2 g^2} \left(\frac{f''}{f} - \frac{f'g'}{fg} + \frac{2f'}{rf} + \frac{2g'}{rg} + \frac{g^2 - 1}{r^2} \right) = YE^2 + \rho, \quad (24)$$

$$\frac{1}{\kappa^2 g^2} \left(\frac{f''}{f} - \frac{f'g'}{fg} - \frac{2f'}{rf} - \frac{2g'}{rg} + \frac{g^2 - 1}{r^2} \right) = YE^2 - \rho/3, \quad (25)$$

$$\frac{1}{\kappa^2 g^2} \left(\frac{f''}{f} - \frac{f'g'}{fg} + \frac{g^2 - 1}{r^2} \right) = YE^2 + \rho/3, \quad (26)$$

and we have the following conservation relation from the covariant exterior derivative of the gravitational field equations (16)

$$p' + 4p \frac{f'}{f} = 2(YE)'E + \frac{4YE^2}{r}, \quad (27)$$

together with the constraint from (17)

$$\frac{dY}{dR} = \frac{1}{2\kappa^2 E^2} \quad (28)$$

where the curvature scalar is

$$R = \frac{2}{g^2} \left(-\frac{f''}{f} + \frac{f'g'}{fg} - \frac{2f'}{fr} + \frac{2g'}{gr} + \frac{g^2 - 1}{r^2} \right). \quad (29)$$

A. EXACT SOLUTIONS UNDER CONFORMAL SYMMETRY

We assume that the existence of a one parameter group of conformal motions for the metric (19)

$$L_\xi g_{ab} = \phi(r)g_{ab} \quad (30)$$

where $L_\xi g_{ab}$ is Lie derivative of the interior metric with respect to the vector field ξ and $\phi(r)$ is an arbitrary function of r . The interior gravitational field of stars can be described by using this symmetry [19],[59–61]. The metric functions $f^2(r)$ and $g^2(r)$ satisfying this symmetry were obtained as

$$f^2(r) = a^2 r^2, \quad g^2(r) = \frac{\phi_0^2}{\phi^2} \quad (31)$$

in [59] where a and ϕ_0 are integration constants. Introducing a new variable $X = \frac{\phi^2}{\phi_0^2}$ in (31) and using this symmetry, equations (24)-(28) turn out to be the three differential equations

$$-\frac{X'}{\kappa^2 r} + \frac{2X + 2}{\kappa^2 r^2} - 2\rho = 2YE^2, \quad (32)$$

$$\frac{X'}{\kappa^2 r} - \frac{2X}{\kappa^2 r^2} + \frac{2\rho}{3} = 0, \quad (33)$$

$$p' + \frac{4p}{r} - 2(YE)'E - \frac{4YE^2}{r} = 0. \quad (34)$$

Here we note that the constraint (28) is not an independent equation from (32) and (33), since we find the constraint eliminating ρ from (32)-(33) and taking derivative of the result equation as in [35] (where $\rho = 3p$). Thus, we have three differential equations (32-34) and four unknowns (X, ρ, Y, E) . So a given theory or a non-minimal coupling function $Y(R)$, it may be possible to find the corresponding exact solutions for the functions X, E and ρ , or inversely, for a convenient choice of any one of the functions X, E and ρ , we may find the corresponding non-minimal theory via the non-minimal function $Y(R)$. In this paper we will continue with the second case offering physically acceptable metric solutions. In the second case, one of the challenging problems is to solve r from $R(r)$ and re-express the function Y depending on R .

When we choose the metric function $g^2(r) = \frac{1}{X} = \frac{3}{1-br^2}$ as a result in [19] with a constant b , we find the constant curvature scalar $R = 4b$ and a constant non-minimal function $Y(R)$. Then this model (1) turns out that the minimal Einstein-Maxwell case. Therefore we need non-constant curvature scalar to obtain non-trivial solutions of the non-minimal theory. Inspired by [19], for $\alpha > 2$ real numbers and $b \neq 0$, we offer the following metric function

$$g^2(r) = \frac{1}{X} = \frac{3}{1+br^\alpha} \quad (35)$$

which is regular at origin $r = 0$ and giving the following non-constant and regular curvature scalar

$$R = -b(\alpha + 2)r^{\alpha-2}. \quad (36)$$

We note that if $b = 0$ the curvature scalar R becomes zero and Y is a constant again. Therefore, here we consider the case with $b \neq 0$ and obtain the following solutions to the equations (32-35)

$$\rho(r) = \frac{2 - br^\alpha(\alpha - 2)}{2\kappa^2 r^2}, \quad (37)$$

$$Y(r) = c [1 + b(\alpha - 2)r^\alpha]^{-\frac{3(\alpha+2)}{2\alpha}}, \quad (38)$$

$$E^2(r) = \frac{[1 + b(\alpha - 2)r^\alpha]^{\frac{5\alpha+6}{2\alpha}}}{3c\kappa^2 r^2}. \quad (39)$$

Here c is a non-zero integration constant and it will be determined by the exterior Einstein-Maxwell Lagrangian (48) as $c = 1$. Using the charge-radius relation (23), we calculate the

total charge inside the volume with radius r

$$q^2(r) = (YEr^2)^2 = \frac{cr^2 [1 + b(\alpha - 2)r^\alpha]^{-\frac{\alpha+6}{2\alpha}}}{3\kappa^2}. \quad (40)$$

We see that the charge is regular at the origin $r = 0$ for the theory with $\alpha > 2$. Obtaining by the inverse of $R(r)$ from (36)

$$r = \left(\frac{-R}{\alpha b + 2b} \right)^{1/(\alpha-2)} \quad (41)$$

the non-minimal coupling function is calculated as

$$Y(R) = c \left[1 + b(\alpha - 2) \left(\frac{-R}{\alpha b + 2b} \right)^{\frac{\alpha}{\alpha-2}} \right]^{-\frac{3\alpha+6}{2\alpha}}. \quad (42)$$

The non-minimal function (42) turns to $Y(R) = c$ for the vacuum case $R = 0$ and we can choose $c = 1$ to obtain the known minimal Einstein-Maxwell theory at the exterior region. Thus the Lagrangian of our non-minimal gravitational theory (1)

$$L = \frac{1}{2\kappa^2} R * 1 - \left[1 + b(\alpha - 2) \left(\frac{-R}{\alpha b + 2b} \right)^{\frac{\alpha}{\alpha-2}} \right]^{\frac{-3(\alpha+2)}{2\alpha}} F \wedge *F + 2A \wedge J + L_{mat} + \lambda_a \wedge T^a \quad (43)$$

admits the following metric

$$ds^2 = -a^2 r^2 dt^2 + \frac{3}{1 + br^\alpha} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \quad (44)$$

together with the energy density, electric field and electric charge

$$\rho(r) = \frac{2 - br^\alpha(\alpha - 2)}{2\kappa^2 r^2}, \quad (45)$$

$$E^2(r) = \frac{[1 + b(\alpha - 2)r^\alpha]^{\frac{5\alpha+6}{2\alpha}}}{3\kappa^2 r^2}, \quad (46)$$

$$q^2(r) = \frac{r^2 [1 + b(\alpha - 2)r^\alpha]^{-\frac{\alpha+6}{2\alpha}}}{3\kappa^2} \quad (47)$$

under the conformal symmetry (30) describing the interior of the radiation fluid star with $\alpha > 2$. The parameter b in the model will be determined from the matching condition (62) and the parameter α can be determined by the related observations.

On the other hand, since the exterior region does not have any matter and source the above non-minimal Lagrangian (43) turns to the following sourceless minimal Einstein-Maxwell Lagrangian,

$$L = \frac{1}{2\kappa^2} R * 1 - F \wedge *F + \lambda_a \wedge T^a, \quad (48)$$

which is the vacuum case with $Y(R) = 1$ and the field equations of the non-minimal theory (2-4) turn to the following Einstein-Maxwell field equations due to $Y_R = 0$

$$d * F = 0, \quad dF = 0, \quad (49)$$

$$-\frac{1}{2\kappa^2} R_{bc} \wedge *e^{abc} = \iota^a F \wedge *F - F \wedge \iota^a *F \quad (50)$$

which lead to $R = 0$ from trace equation and admit the following Reissner-Nordstrom metric

$$ds^2 = -\left(1 - \frac{2M}{r} + \frac{\kappa^2 Q^2}{r^2}\right) dt^2 + \left(1 - \frac{2M}{r} + \frac{\kappa^2 Q^2}{r^2}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (51)$$

with the electric field

$$E(r) = \frac{Q}{r^2} \quad (52)$$

at the exterior region. Here M is the total gravitational mass and $Q = q(r_b)$ is the total charge of the star. Since the Ricci scalar is zero for the Reissner-Nordstrom solution, the non-minimal function (42) becomes $Y = 1$ consistent with the above considerations. As we see from (49) the excitation 2-form $\mathcal{G} = YF$ is replaced by the Maxwell tensor F at the exterior vacuum region. In order to see a concrete example of this non-minimally coupled theory we look on the simplest case where $\alpha = 3$, then the non-minimal Lagrangian is

$$L = \frac{1}{2\kappa^2} R * 1 - \left(1 - \frac{R^3}{5^3 b^2}\right)^{-\frac{5}{2}} F \wedge *F + 2A \wedge J + L_{mat} + \lambda_a \wedge T^a \quad (53)$$

and its corresponding field equations admit the following interior metric

$$ds^2 = -a^2 r^2 dt^2 + \frac{3}{1 + br^3} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (54)$$

Using the curvature scalar $R = -5br$, we find the energy density, electric field and charge as

$$\rho(r) = \frac{1}{\kappa^2 r^2} - \frac{b}{2\kappa^2} r, \quad (55)$$

$$E^2(r) = \frac{(1 + br^3)^{\frac{7}{2}}}{3\kappa^2 r^2}, \quad (56)$$

$$q(r) = \frac{r^2(1 + br^3)^{-3/2}}{3\kappa^2}. \quad (57)$$

For the exterior region ($R = 0$), the model (53) turns to the well known Einstein-Maxwell theory which admits the above Reissner-Nordstrom solution.

IV. MATCHING CONDITIONS

We will match the interior and exterior metric (44), (51) at the boundary of the star $r = r_b$ for continuity of the gravitational potential

$$a^2 r_b^2 = 1 - \frac{2M}{r_b} + \frac{\kappa^2 Q^2}{r_b^2}, \quad (58)$$

$$\frac{3}{1 + br_b^\alpha} = \left(1 - \frac{2M}{r_b} + \frac{\kappa^2 Q^2}{r_b^2}\right)^{-1}. \quad (59)$$

The matching conditions (58) and (59) give

$$a^2 = \frac{\kappa^2 Q^2 - 2Mr_b + rb^2}{r_b^4}, \quad (60)$$

$$b = \frac{2r_b^2 - 6Mr_b + 3\kappa^2 Q^2}{r_b^{2+\alpha}}. \quad (61)$$

The vanishing pressure condition at the boundary r_b requires that

$$p(r_b) = \frac{2 - b(\alpha - 2)r_b^\alpha}{6\kappa^2 r_b^2} = 0, \quad (62)$$

and it determines the constant b in the non-minimal model (43) as

$$b = \frac{2}{(\alpha - 2)r_b^\alpha}. \quad (63)$$

The interior region of the star can be considered as a specific medium and the exterior region as a vacuum. Then the excitation 2-form $\mathcal{G} = YF$ in the interior turns to the Maxwell tensor F at the exterior, because of $Y = 1$ in this vacuum region. That is, we use the continuity of the tensor at the boundary which leads to the continuity of the total charge in which a volume V . Then the total charge for the exterior region is obtained from the Maxwell equation (49), $d * F = 0$, taking the integral $\frac{1}{4\pi} \int_{\partial V} * F = Er^2 = Q$, while the total charge in the interior region is given by (23). Thus the total charge Q is determined by setting $r = r_b$ in (47) as a last matching condition

$$q^2(r_b) = \frac{r_b^2 [1 + b(\alpha - 2)r_b^\alpha]^{-\frac{\alpha+6}{2\alpha}}}{3\kappa^2} = Q^2 . \quad (64)$$

Substituting (63) in (64) we obtain the following total charge-boundary radius relation

$$Q^2 = \frac{r_b^2}{\kappa^2 3^{\frac{3\alpha+6}{2\alpha}}} . \quad (65)$$

The ratio $\frac{\kappa^2 Q^2}{r_b^2} = 3^{-\frac{3\alpha+6}{2\alpha}}$ which is obtained from (65) is plotted in Figure 1 depending on the parameter α of the model for different α intervals. As we see from (65) the charge-radius ratio has the upper limit

$$\frac{\kappa^2 Q^2}{r_b^2} = \frac{1}{3\sqrt{3}} \approx 0.1924 . \quad (66)$$

When we compare (63) with (61), we find the following mass-charge relation for the model with the non-minimally coupled electromagnetic fields to gravity

$$M = \left(\frac{\alpha - 3}{\alpha - 2}\right) \frac{r_b}{3} + \frac{\kappa^2 Q^2}{2r_b} . \quad (67)$$

Substituting the total charge (65) in (67) we find the total mass of the star depending on the boundary radius r_b and the parameter α of the model

$$M = \left(\frac{\alpha - 3}{\alpha - 2}\right) \frac{r_b}{3} + 3^{-\frac{3\alpha+6}{2\alpha}} \frac{r_b}{2} . \quad (68)$$

This mass-radius relation is shown in Figure 2 for two different α intervals. Taking the limit $\alpha \rightarrow \infty$, we can find the upper bound for the mass-radius ratio

$$\frac{M}{r_b} < \frac{1}{3} + \frac{1}{6\sqrt{3}} \approx 0.4295 \quad (69)$$




FIG. 1: The square of the charge-radius ratio versus the parameter α .

which is slightly smaller than Buchdahl bound [44] and the bound given in [43] for General Relativistic charged objects. Also, the matter mass component of the radiation fluid star is




FIG. 2: The gravitational mass-radius ratio versus the parameter α .

obtained from the following integral of the energy density ρ

$$M_m = \frac{\kappa^2}{2} \int_0^{r_b} \rho r^2 dr = \frac{\alpha}{2(\alpha + 1)} r_b. \quad (70)$$

The upper bound of the matter mass for the radiative star is found as $M_m = \frac{r_b}{2}$ taking by the limit $\alpha \rightarrow \infty$. The dependence of the matter mass-radius ratio on the parameter α can be seen in Figure 3. Here we emphasize that each different value of α corresponds to a




FIG. 3: The matter mass-radius ratio versus the parameter α .

different non-minimally coupled theory in (43) and the each different theory gives a different mass-radius relation. Additionally, the gravitational surface redshift z is calculated from

$$z = \left(1 - \frac{2M}{r_b} + \frac{\kappa^2 Q^2}{r_b^2}\right)^{-\frac{1}{2}} - 1 = \sqrt{\frac{3(\alpha - 2)}{\alpha}} - 1. \quad (71)$$

Taking the limit $\alpha \rightarrow \infty$, the inequality for the redshift is found as $z < \sqrt{3} - 1 \approx 0.732$ which is smaller than the bound given in [43] and the Buchdahl bound $z \leq 2$. We plot the redshift depends on the α in Fig. 4.

For the case $\alpha = 3$, we calculate all the parameters as $M = \frac{\sqrt{3}r_b}{54} \approx 0.032r_b$, $M_m = \frac{3r_b}{8} = 0.375r_b$, $Q^2 = \frac{\sqrt{3}r_b^2}{27\kappa^2} \approx \frac{0.064r_b^2}{\kappa^2}$ from (68), (70), (65). In this case, because of $\frac{2M}{r_b} = \frac{\kappa^2 Q^2}{r_b^2}$ at the boundary, the metric functions are equal to 1, $f(r_b) = g(r_b) = 1$. This means that the total gravitational mass M together with the energy of the electromagnetic field inside the boundary is exactly balanced by the energy of the electromagnetic field outside the boundary and then the gravitational surface redshift becomes zero from (71) for this case $\alpha = 3$. In Table 1, we determine some α values in the model for some specific mass-radius relations




FIG. 4: The gravitational surface redshift versus the parameter α .

Star	α	$\frac{M}{r_b}$	$M(M_\odot)$	$r_b(km)$	$\frac{\kappa^2 Q^2}{r_b^2}$	z (redshift)
SMC X-1	3.453	0.141	1.29	9.13	0.074	0.124
Cen X-3	3.555	0.157	1.49	9.51	0.076	0.145
PSR J1903+327	3.648	0.170	1.667	9.82	0.078	0.164
Vela X-1	3.702	0.177	1.77	9.99	0.079	0.174
PSR J1614-2230	3.822	0.191	1.97	10.3	0.081	0.195

TABLE I: Some values of α and the corresponding other parameters for some given mass and mass-radius relations of neutron stars

in the literature. As we see from the relation (68) each α gives a mass-radius ratio. Then taking also the observed mass values of some neutron stars from the literature we can find the corresponding values of the parameters such as the boundary radius, charge-radius ratio and redshift.

V. CONCLUSION

We have analyzed the exact solutions of the non-minimally coupled $Y(R)F^2$ theory for the radiation fluid stars which have the equation of state $\rho = 3p$, assuming the existence of a one parameter group of conformal motions. We have found new solutions which lead to regular metric functions and regular Ricci scalar inside the star. We have obtained non-negative matter density ρ and pressure p which vanish at the boundary of the star $r = r_b$, $\rho = 3p = \frac{r_b^\alpha - r^\alpha}{\kappa^2 r_b^2 r^2}$. The derivatives of the density and pressure are negative as required for acceptable interior solution, that is, $\frac{d\rho}{dr} = 3\frac{dp}{dr} = -\frac{(\alpha-2)r^\alpha + 2r_b^\alpha}{\kappa^2 r_b^\alpha r^3}$ (where $\alpha > 2$). The speed of sound $(\frac{dp}{d\rho})^{1/2} = \frac{1}{\sqrt{3}} < 1$ satisfies the implication of causality, since it does not exceed the speed of light $c = 1$. But the mass density ρ and charge density ρ_e have singularity at the center of the star as the same feature in [19]. However, this feature is physically acceptable since the total charge and mass became finite for the model.

After obtaining the exterior and interior metric solutions of the non-minimal theory, we matched them at the boundary r_b . Using the vanishing pressure condition and total charge at the boundary, we obtained the square of the total charge-radius ratio $\frac{\kappa^2 Q^2}{r_b^2}$, mass-radius ratio $\frac{M}{r_b}$ and gravitational surface redshift z depending on the parameter α of the model. Taking the limit $\alpha \rightarrow \infty$, we found the ratio $\frac{\kappa^2 Q^2}{r_b^2}$ which has the upper bound $\frac{1}{3\sqrt{3}} \approx 0.1924$ and the mass-radius ratio which has the upper bound $\frac{M}{r_b} = \frac{1}{3} + \frac{1}{6\sqrt{3}} \approx 0.4295$. We note that this maximum mass-radius ratio is smaller than the bound which was found by Mak *et al.* [43] for charged General Relativistic objects even also Buchdahl bound $4/9$ [44] for uncharged compact objects. Also we found the upper limit $z = \sqrt{3} - 1 \approx 0.732$ for the gravitational surface redshift in the non-minimal model and it satisfies the bound given in [43] for charged stars. On the other hand the minimum redshift $z = 0$ corresponds to the parameter $\alpha = 3$. We have plotted all these quantities depend on the parameter α .

We determined some values of the parameter α in Table 1 for some specific mass-radius relations which were given by the literature. Also using the observed mass values we found the corresponding parameters such as the boundary radius, charge-radius ratio and redshift for some known neutron stars. It would be interesting to generalize the analysis to the extended theories of gravity [20, 62] coupled to the Maxwell theory in future studies.

ACKNOWLEDGEMENT

I would like to thank to the anonymous referee for very useful comments and suggestions.

- [1] J.R. Oppenheimer and G.M. Volkoff, *Phys. Rev.* **55**, 374 (1939).
- [2] C. W. Misner and H. S. Zepolsky, *Phys. Rev. Lett.*, **12**, 635 (1964).
- [3] D.W. Meltzer, K.S. Thorne, *Astrophys. J.*, **145**, 514, (1966).
- [4] S. Chandrasekhar, *An Introduction to the Theory of Stellar Structure* (Dover), (1942).
- [5] R.D. Sorkin, R.M. Wald and Z. Zhen Jiu, *Gen. Rel. Grav.* 13 1127, (1981).
- [6] H.-J. Schmidt and F. Homann, *Gen. Rel. Grav.* 32, 919 (2000).
- [7] P. H. Chavanis, *Astron. Astrophys.* 483, 673 (2008).
- [8] P.H. Chavanis, *Astron. Astrophys.* 381, 709, (2002).
- [9] A. Mitra and N. K. Glendenning, *Monthly Notices of the Royal Astronomical Society: Letters* 404, L50 (2010).
- [10] N. K. Glendenning, *Compact Stars, Nuclear Physics, Particle Physics and General Relativity*, Springer, New York, (2000).
- [11] Penrose, R. (1979). In *General Relativity: An Einstein Centenary Survey*, eds. S. W. Hawking and W. Israel, Cambridge University Press (1981).
- [12] R. M. Wald, *Phys. Rev. D*, 21, 2742, (1980).
- [13] M. K. Mak, and T. Harko, *Pramana*, 65, 185, (2005).
- [14] M. K. Mak, and T. Harko, *The European Physical Journal C*, 73, 2585 (2013).
- [15] T. Harko, M. K. Mak, *Astrophysics and Space Science*, 361(9), 1-19 (2016).
- [16] R. Stettner, *Ann. Phys.* 80, 212 (1973).
- [17] A. Krasinski, *Inhomogeneous cosmological models*, Cambridge University Press (1997).
- [18] R. Sharma, S. Mukherjee and S.D. Maharaj, *Gen. Relativ. Gravit.* 33, 999 (2001).
- [19] M. K. Mak, and T. Harko, *Int. J. Mod. Phys. D*, 13, 149 (2004).
- [20] B. Danila, T. Harko, F. S. N. Lobo, M. K. Mak, "Hybrid metric-Palatini stars", arXiv:1608.02783 [gr-qc].
- [21] T. Harko, F. S. N. Lobo, M. K. Mak, and S. V. Sushkov, *Phys. Rev. D* 88, 044032 (2013).

- [22] A. R. Prasanna, Phys. Lett. 37A, 331 (1971).
- [23] G. W. Horndeski, J. Math. Phys. (N.Y.) 17, 1980 (1976).
- [24] F. Mueller-Hoissen, Class. Quant. Grav. 5, L35 (1988)
- [25] T. Dereli, G. Üçoluk, Class. Q. Grav. **7**, 1109 (1990).
- [26] H.A. Buchdahl, J. Phys. A 12, 1037 (1979)
- [27] I.T. Drummond, S.J. Hathrell, Phys. Rev. D 22, 343 (1980)
- [28] M. S. Turner and L. M. Widrow, Phys. Rev. D 37, 2743 (1988).
- [29] F. D. Mazzitelli and F. M. Spedalieri, Phys. Rev. D 52, 6694 (1995) [astro-ph/9505140].
- [30] L. Campanelli, P. Cea, G. L. Fogli and L. Tedesco, Phys. Rev. D 77, 123002 (2008) [arXiv:0802.2630 [astro-ph]].
- [31] K. E. Kunze, Phys. Rev. D 81, 043526 (2010) [arXiv:0911.1101 [astro-ph.CO]].
- [32] K. Bamba and S. D. Odintsov, JCAP (04), 024 (2008).
- [33] T. Dereli, Ö. Sert, Mod. Phys. Lett. A **26**, 20, 1487-1494 (2011).
- [34] A. Baykal, T. Dereli, Phys.Rev. D92 6, 065018 (2015).
- [35] Ö. Sert, Journal of Mathematical Physics **57**, 032501 (2016).
- [36] T. Dereli, Ö. Sert, Eur. Phys. J. C **71**, 3, 1589 (2011).
- [37] Ö. Sert, Eur. Phys. J. Plus 127: 152 (2012).
- [38] Ö. Sert, Mod. Phys. Lett. A **28**, 12, 1350049 (2013).
- [39] Ö. Sert and M. Adak, arXiv:1203.1531 [gr-qc].
- [40] M. Adak, Ö. Akarsu, T. Dereli, Ö. Sert, "Anisotropic inflation with a non-minimally coupled electromagnetic field to gravity", arXiv:1611.03393 [gr-qc].
- [41] K. Bamba, S. Nojiri and S. D. Odintsov, JCAP, (10), 045, (2008).
- [42] T. Dereli, Ö. Sert, Phys. Rev. D **83**, 065005 (2011).
- [43] M. K. Mak, Peter N. Dobson Jr., T. Harko Europhys.Lett. 55 310-316 (2001).
- [44] H.A. Buchdahl, Phys. Rev., 116 1027 (1959).
- [45] J. M. Overduin and P. S. Wesson, Physics Reports 402, 267 (2004).
- [46] H. Baer, K.-Y. Choi, J. E. Kim, and L. Roszkowski, Physics Reports 555, 1 (2015).
- [47] A. G. Riess et al., Astron. J. 116, 1009 (1998).
- [48] S. Perlmutter et al., Astrophys. J. 517, 565 (1999).
- [49] R. A. Knop et al., Astrophys. J. 598, 102 (2003).
- [50] R. Amanullah et al., Astrophys. J. 716, 712 (2010).

- [51] D. H. Weinberg, M. J. Mortonson, D. J. Eisenstein, C. Hirata, A. G. Riess, and E. Rozo, Physics Reports 530, 87 (2013).
- [52] D. J. Schwarz, C. J. Copi, D. Huterer and G. D. Starkman, Class. Quant. Grav. 33 184001 (2016).
- [53] T. Dereli, J. Gratus, R.W. Tucker, J. Phys. A, Math. Theor. 40, 5695 (2007).
- [54] T. Dereli, J. Gratus, R.W. Tucker, Phys. Lett. A 361(3), 190193 (2007).
- [55] A. Einstein, Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.) 1919, 433 (1919).
- [56] S. Weinberg, Rev. Mod. Phys. 61, 1 (1989).
- [57] W. G. Unruh, Phys. Rev. D 40, 1048 (1989).
- [58] G. F. R. Ellis, H. van Elst, J. Murugan and J. P. Uzan, Class. Quant. Grav. 28, 225007 (2011).
- [59] L. Herrera and J. Ponce de Leon, J. Math. Phys. 26, 2303, (1985).
- [60] L. Herrera and J. Ponce de Leon, J. Math. Phys. 26, 2018, (1985).
- [61] L. Herrera and J. Ponce de Leon, J. Math. Phys. 26, 778, (1985).
- [62] A. V. Astashenok, S. Capozziello, S. D. Odintsov, JCAP 01 001 (2015).